

Thames Valley Priorities Committee Commissioning Policy Statement

Policy No. TVPC34 **Erectile Dysfunction**

**Recommendation made by
the Priorities Committee:** **November 2015; reviewed January 2019¹**

Date of issue (v3.0): **November 2021²**

Date of issue (v2.1): **January 2019**

Date of issue (v2.0): **July 2017**

Date of issue (v1.0): **July 2016**

The Thames valley Priorities Committee has considered the evidence of clinical and cost effectiveness of treatments for erectile dysfunction in adult patients. The publication of the updated Statutory Instrument 2014/1625¹ was taken into account in making the following recommendations:

1. Funding for treatment with the phosphodiesterase type -5 inhibitor, generic sildenafil, at the minimum effective dose is **recommended** for any man presenting with erectile dysfunction with a frequency of dosing of four times per month.
2. Funding for all other phosphodiesterase type-5 inhibitors (vardenafil, tadalafil and avanafil) is only recommended for patients who meet the Government Selected List Scheme (SLS) criteria^{1,2} AND where generic sildenafil is ineffective, with a frequency of dosing of two times per month using the drug with the lowest acquisition cost.
3. Funding for treatment with prostaglandin E1 intracavernosal injections and intraurethral instillations is only recommended for patients who meet the SLS criteria^{1,2} AND only if oral phosphodiesterase type-5 inhibitors are contraindicated or ineffective. The maximum frequency of dosing should be two times per month using the drug with the lowest acquisition cost.
4. Treatment with Alprostadil cream is **not normally funded** in view of limited evidence for clinical and cost effectiveness.
5. Treatment with vacuum erection devices is **not normally funded** in view of limited evidence for clinical and cost effectiveness.
6. Treatment with penile implants is not normally funded by the CCGs. Penile Prosthesis surgery for end stage erectile dysfunction is commissioned by NHS England, please see [NHS England Clinical Commissioning policy \(2016\)](#)
7. Treatment with psychosexual interventions is **not normally funded** in view of limited evidence for clinical and cost effectiveness.

Phosphodiesterase type-5 inhibitors (sildenafil, vardenafil and tadalafil) are oral drugs that enable a penile erection with sexual stimulation. There is evidence for the effectiveness of these drugs in men with erectile dysfunction of varying causes. The effectiveness of individual drugs is comparable and generic sildenafil has been shown to be cost-effective. The SLS restrictions set

¹ No changes have been made to this policy.

² Addition of audit codes

out in Statutory Instrument 1999/1627² apply to all phosphodiesterase type-5 inhibitors except generic sildenafil.

Prostaglandin E1 is used in the treatment of erectile dysfunction when delivered locally into penile tissues. There is evidence for the effectiveness of intra-cavernosal injections of prostaglandin E1 in those unresponsive to oral drugs.

Psychosexual interventions such as counselling and psychotherapy comprise a group of techniques with limited evidence for effectiveness and no evidence of cost-effectiveness.

Vacuum erection devices are mechanical devices for producing an erection that is sustained with the placement of a constricting band across the base of the penis. There is limited evidence for the effectiveness of these devices and no evidence of cost-effectiveness.

Penile implants can be surgically inserted into the penis for treating erectile dysfunction. There is limited evidence of effectiveness with a high proportion of patients having major complications.

Audit codes

ICD10 diagnosis code:

F52.2 Failure of genital response

OPCS procedure codes relevant to erectile dysfunction in conjunction with the above ICD10 code:

N29.1 Implantation of prosthesis into penis

N32.4 Injection of therapeutic substance into penis [includes penile injection to produce erection]

N32.6 Operations on penis for erectile dysfunction NEC

NOTES:

- Potentially exceptional circumstances may be considered by a patient's CCG where there is evidence of significant health status impairment (e.g. inability to perform activities of daily living) and there is evidence that the intervention sought would improve the individual's health status.
- This policy will be reviewed in the light of new evidence or new national guidance, e.g. from NICE.
- Thames Valley clinical policies can be viewed at <http://www.fundingrequests.cscsu.nhs.uk/>

¹ http://www.legislation.gov.uk/uksi/2014/1625/pdfs/uksi_20141625_en.pdf

² http://www.legislation.gov.uk/uksi/1999/1627/pdfs/uksi_19991627_en.pdf